

Health Services DHS Pediatrics (0-18) Asthma Action Plan www.labreathmobile.com

Patient Name		e:	MRN:	Date of Birth:	
Parent Name:			D.		
Clinic/PCP:			51	Child's next appointment:	
School:			Phone:		
	þo	 Breathing is good No cough or wheeze Can work and play 	PREVENT asthma symptoms every day: (A DHS Formulary Medicine: How much QVAR® 40, 80mcg Puffs Flovent® 44¹, 110¹, 220²mcg Puffs Puffs	n: When: times per day	
GREEN	l Feel Good		□ PULMICORT®³ 90, 180mcg Puffs □ Advair®⁴ 100/50, 250/50mcg Puffs □ Dulera®⁵ 100/5, 200/5mcg Puffs □ Montelukast 4, 5, 10mg Table	times per day times per day ts times per day	
		Peak Flow Number	Budesonide® 0.25, 0.5mg in nebulizer	times per day	
		to		ricted to pregnancy; ⁴ Restricted to tricted to children<9	
		 Cough or wheeze Difficulty breathing SLOW DOWN & take relief medicine: (Rinse mouth after using inhalers) 			
		Wake up at night	DHS Formulary Medicine: How muc	<u>:h:</u> When:	
	<u>I</u> Feel Good			uffs times per day	
			Albuterol/*levalbuterol in nebul		
YELLOW				 ' '	
		Cough Wheeze	(*only if on Managed Care formulary) ALSO CONTINUE/INCREASE your preventive medicine:		
▎≓				Puffs times per day	
}	NOT			Puffs times per day	
		Tight Wake up chest at night		Puffs times per day	
	Do	· ,		uffs times per day	
	_	Peak Flow Number		ouffs times per day	
		to (50-80% of personal best)	Montelukast [®] 4, 5, 10mg T		
		(30-80% of personal best)	☐ Budesonide ^{®6} 0.25, 0.5mg in nebu	lizer times per day	
		Medicine not helping MEDICAL ALERT – GET HELP NOW! Call your doctor at			
		Breathing hard, fastCan't talk/walk well	eathing hard, fast Take these medicines until you talk to the doctor or for school, until you talk to the parent:		
			DHS Formulary Medicine: How n	nuch: When:	
	wful	Repeat albuterol/	Albuterol/*levalbuterol in nebuterol (*only if on Managed	' '	
RED	Feel <u>Awful</u>	levalbuterol in 20 minutes if needed x 3	Prednisone (1-2mg/kg/day) Tablets as follows:		
	H	inceded A 3	Prednisolone 15mg/5cc; 5mg/5cc	Teaspoons as follows:	
		Peak Flow Number	Continue to use all medications in the	e yellow zone - shaded box	
		to (<50% of personal best)	Don't wait – Call 911 if your asthma is severe	or if no improvement after medicine	
Authorization and Disclaimer from Parent/Guardian: Action Plan in accordance with state laws and regulations. I request that the school assist my child with the above asthma medications and the Asthma Yes No					
My child may carry and self-administer asthma medications and I agree to release the school district and school personnel from all claims of liability if my child suffers any adverse reactions from self-administration of asthma medications.					
Print Parent/Guardian Name: Signature: Date:					
<u>Health Care Provider:</u> My signature provides authorization for the above written orders. I understand that all procedures will be implemented in accordance with state laws and regulations. Student may carry and self-administer asthma medications: Yes No					
(This authorization is for a maximum of one year from sig Print Provider Name/Credentials:				Date: v11 NU-091613	

This guide suggests actions that you can do to avoid your asthma triggers (anyone with asthma should have an Asthma Action Plan)

(Check Mark is Next to Your Triggers)





Animal Dander

Some people are allergic to the flakes of skin or dried saliva from animals with fur or feathers. The best thing to do is not have a pet if you are allergic.

If you cannot keep the pet outdoors, then:

- Keep the pet out of your bedroom, and keep the door closed. Keep pets away from fabric-covered furniture and carpets.
- Remove carpets and furniture covered with cloth from your home.
- Air purifiers with HEPA-filter may reduce dander.



_Dust Mites

Dust mites are tiny bugs that are found in mattresses, pillows, carpets, upholstered furniture, bedcovers, clothes, stuffed toys, and fabric or other fabric-covered items.

- Cover your mattress in special dust-mite proof encasings.
- Cover your pillow in a special dust-mite proof encasing or wash the pillow each week in hot water. Water must be hotter than 130°F to kill the mites
- Wash the sheets and blankets on your bed each week in hot water.
- Reduce indoor humidity to below 60% (ideally between 30-50%). Dehumidifiers or central air conditioners can do this.
- Try not to sleep or lie on cloth-covered cushions.
- Remove carpets from your bedroom and those laid on concrete, if possible. Vacuum twice weekly with HEPA-filter vacuum.
- Keep stuffed toys out of the bed or wash weekly in hot water.



Cockroaches

Many people with asthma are allergic to the dried droppings and remains of cockroaches.

- Keep food and garbage in closed containers.
 Never leave food out.
- Use poison baits, powder, gels, traps, or paste (for example, boric acid). Heavy infestations may require professional treatment. (Stay out of the room until the odor goes away if spray used).



30

___Indoor Mold

- Fix leaky faucets, pipes, or other sources of water that have mold around them.
- Clean moldy surfaces with a cleaner that has bleach in it.
- Extensive indoor mold may require professional assistance.

(Stamp Plate)





__Pollen and Outdoor Mold **

What to do during your allergy season (when pollen or mold spore counts are high):

- Try to keep your windows and doors closed (especially in the early morning for tree allergy)
- Stay indoors with windows closed from late morning to afternoon when pollen and mold spore counts are high.
- Ask your doctor whether you need to take or increase allergy and/or anti-inflammatory medicines before your allergy season starts.





Irritants:

Tobacco Smoke

- Avoid being around any type of smoke.
- If you smoke, ask your doctor for ways to help you quit. Ask family members to quit smoking.
- Do not allow smoking in the home or car.
- If someone smokes, ask them to change their clothes and wash their hands before coming close to you.

__Smoke, Strong Odors, and Sprays

- If possible, do not use a wood-burning stove, kerosene heater, or fireplace.
- Try to stay away from strong odors and sprays, such as perfume, talcum powder, hair spray, and paints.

Other things that can make asthma worse:

Vacuum Cleaning

- Try to have someone else vacuum for you once or twice a week. Stay out of the room while the room is being vacuumed and for at least 20 minutes afterward.
- If you vacuum, use a dust mask (from hardware store), a double-layerd or microfilter vacuum cleaner bag, or a vacuum cleaner with a HEPA filter.

____Cold Air: Cover your nose and mouth with a scarf on cold or windy days.

____ Food Allergies: Avoid any food which you have been told is a food allergy.

_____ Respiratory Virus and Cold Symptoms

Talk to your provider about your asthma action
plan and using scheduled rescue medications at the
onset of respiratory symptoms. See you provider
for annual flu vaccine.

